

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 May 2001 (10.05.2001)

PCT

(10) International Publication Number
WO 01/32209 A1

(51) International Patent Classification⁷: **A61K 39/395**,
39/40, 39/42, 39/44, 39/00, 39/38, 38/00, C12N 5/06, 5/16

Sei-Yu; 160 Mira Street, Foster City, CA 94404 (US). SUN,
Yongming; Apartment 260, 869 S. Winchester Boulevard,
San Jose, CA 95128 (US).

(21) International Application Number: PCT/US00/30482

(74) Agents: LICATA, Jane, Massey et al.; Licata & Tyrrell
P.C., 66 E. Main Street, Marlton, NJ 08053 (US).

(22) International Filing Date:
3 November 2000 (03.11.2000)

(81) Designated States (*national*): CA, JP.

(25) Filing Language: English

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

(26) Publication Language: English

(30) Priority Data:
60/163,444 4 November 1999 (04.11.1999) US

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

(71) Applicant: DIADEXUS, INC. [US/US]; 3303 Octavius
Drive, Santa Clara, CA 95054 (US).

(72) Inventors: RECIPON, Herve; 85 Fortuna Avenue, San
Francisco, CA 94115 (US). MACINA, Roberto, A.; 4118
Crescendo Avenue, San Jose, CA 95136 (US). CHEN,

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: A NOVEL METHOD OF DIAGNOSING, MONITORING, STAGING, IMAGING AND TREATING CANCER

(57) Abstract: The present invention provides new methods for detecting, diagnosing, monitoring, staging, prognosticating, imag-
ing and treating cancer.

WO 01/32209 A1

A NOVEL METHOD OF DIAGNOSING,
MONITORING, STAGING, IMAGING AND TREATING CANCER

FIELD OF THE INVENTION —

This invention relates, in part, to newly developed
5 assays for detecting, diagnosing, monitoring, staging,
prognosticating, imaging and treating cancers, particularly
lung cancer.

BACKGROUND OF THE INVENTION

Lung cancer is the second most prevalent type of
10 cancer for both men and women in the United States and is
the most common cause of cancer death in both sexes. Lung
cancer can result from a primary tumor originating in the
lung or a secondary tumor which has spread from another
organ such as the bowel or breast. Primary lung cancer is
15 divided into three main types; small cell lung cancer; non-
small cell lung cancer; and mesothelioma. Small cell lung
cancer is also called "Oat Cell" lung cancer because the
cancer cells are a distinctive oat shape. There are three
types of non-small cell lung cancer. These are grouped
20 together because they behave in a similar way and respond
to treatment differently to small cell lung cancer. The
three types are squamous cell carcinoma, adenocarcinoma,
and large cell carcinoma. Squamous cell cancer is the most
common type of lung cancer. It develops from the cells
25 that line the airways. Adenocarcinoma also develops from
the cells that line the airways. However, adenocarcinoma
develops from a particular type of cell that produces mucus
(phlegm). Large cell lung cancer has been thus named
because the cells look large and rounded when they are
30 viewed under a microscope. Mesothelioma is a rare type of
cancer which affects the covering of the lung called the
pleura. Mesothelioma is often caused by exposure to
asbestos.

-2-

Secondary lung cancer is cancer that has started somewhere else in the body (for example, the breast or bowel) and spread to the lungs. Choice of treatment for secondary lung cancer depends on where the cancer started.

- 5 In other words, cancer that has spread from the breast should respond to breast cancer treatments and cancer that has spread from the bowel should respond to bowel cancer treatments.

10 The stage of a cancer indicates how far a cancer has spread. Staging is important because treatment is often decided according to the stage of a cancer. The staging is different for non-small cell and for small cell cancers of the lung.

Non-small cell cancer can be divided into four stages.
15 Stage I is very localized cancer with no cancer in the lymph nodes. Stage II cancer has spread to the lymph nodes at the top of the affected lung. Stage III cancer has spread near to where the cancer started. This can be to the chest wall, the covering of the lung (pleura), the middle
20 of the chest (mediastinum) or other lymph nodes. Stage IV cancer has spread to another part of the body.

Since small cell lung cancer can spread quite early in development of the disease, small cell lung cancers are divided into only two groups. These are: limited disease,
25 that is cancer that can only be seen in one lung and in nearby lymph nodes; and extensive disease, that is cancer that has spread outside the lung to the chest or to other parts of the body. Further, even if spreading is not apparent on the scans, it is likely that some cancer cells
30 will have broken away and traveled through the bloodstream or lymph system. To be safe, it is therefore preferred to treat small cell lung cancers as if they have spread, whether or not secondary cancer is visible. Because surgery is not typically used to treat small cell cancer,
35 except in very early cases, the staging is not as critical

-3-

as it is with some other types of cancer. Chemotherapy with or without radiotherapy is often employed. The scans and tests done at first will be used later to see how well a patient is responding to treatment.

5 Procedures used for detecting, diagnosing, monitoring, staging, and prognosticating lung cancer are of critical importance to the outcome of the patient. For example, patients diagnosed with early lung cancer generally have a much greater five-year survival rate as compared to the
10 survival rate for patients diagnosed with distant metastasized lung cancer. New diagnostic methods which are more sensitive and specific for detecting early lung cancer are clearly needed.

 Lung cancer patients are closely monitored following
15 initial therapy and during adjuvant therapy to determine response to therapy and to detect persistent or recurrent disease of metastasis. There is clearly a need for a lung cancer marker which is more sensitive and specific in detecting lung cancer, its recurrence and progression.

20 Another important step in managing lung cancer is to determine the stage of the patient's disease. Stage determination has potential prognostic value and provides criteria for designing optimal therapy. Generally, pathological staging of lung cancer is preferable over
25 clinical staging because the former gives a more accurate prognosis. However, clinical staging would be preferred were it at least as accurate as pathological staging because it does not depend on an invasive procedure to obtain tissue for pathological evaluation. Staging of lung
30 cancer would be improved by detecting new markers in cells, tissues, or bodily fluids which could differentiate between different stages of invasion.

 U.S. Patent 5,877,290 and U.S. Patent 5,837,498, which are incorporated herein by reference, disclose a human
35 Corpuscles of Stannius, staniocalcin polypeptide and the

-4-

nucleic acid sequence encoding this polypeptide. Also disclosed are methods of using this polypeptide for therapeutic purposes such as treatment of electrolyte disorders and disorders due to elevated bone resorption.

5 It has now been found that this polypeptide and the nucleic acid encoding this polypeptide, which are referred to herein as Lngl08 are diagnostic markers for cancer. Accordingly, in the present invention methods are provided for detecting, diagnosing, monitoring, staging,
10 prognosticating, in vivo imaging and treating cancer via Lngl08. Lngl08 refers, among other things, to native proteins expressed by the gene comprising the polynucleotide sequence of SEQ ID NO:1 or 2. The deduced amino acid sequence of a polypeptide encoded thereby is
15 depicted in SEQ ID NO:3. By "Lngl08" it is also meant herein polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1 or 2, but which still encode the same protein. In the alternative, what is meant by Lngl08
20 as used herein, means the native mRNA encoded by the gene comprising SEQ ID NO:1 or 2 or it can refer to the actual gene comprising SEQ ID NO:1 or 2, or levels of a polynucleotide which is capable of hybridizing under stringent conditions to the antisense sequence of SEQ ID
25 NO:1 or 2.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the
30 specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the

-5-

following description and from reading the other parts of the present disclosure.

SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the present invention to provide a method for diagnosing the presence cancer by analyzing for changes in levels of Lng108 in cells, tissues or bodily fluids compared with levels of Lng108 in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein a change in levels of Lng108 in the patient versus the normal human control is associated with cancer.

Further provided is a method of diagnosing metastatic cancer in a patient having cancer which is not known to have metastasized by identifying a human patient suspected of having cancer that has metastasized; analyzing a sample of cells, tissues, or bodily fluid from such patient for Lng108; comparing the Lng108 levels in such cells, tissues, or bodily fluid with levels of Lng108 in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in Lng108 levels in the patient versus the normal human control is associated with cancer which has metastasized.

Also provided by the invention is a method of staging cancer in a human with cancer by identifying a human patient having cancer; analyzing a sample of cells, tissues, or bodily fluid from such patient for Lng108; comparing Lng108 levels in such cells, tissues, or bodily fluid with levels of Lng108 in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in Lng108 levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of Lng108 is associated with a cancer which is regressing or in remission.

-6-

Further provided is a method of monitoring cancer in a human patient for the onset of metastasis. The method comprises identifying a human patient having cancer that is not known to have metastasized; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for Lng108; comparing the Lng108 levels in such cells, tissue, or bodily fluid with levels of Lng108 in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in Lng108 levels in the patient versus the normal human control is associated with a cancer which has metastasized.

Further provided is a method of monitoring the change in stage of cancer in a human patient by looking at levels of Lng108 in the human patient. The method comprises identifying a human patient having cancer; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for Lng108; comparing the Lng108 levels in such cells, tissue, or bodily fluid with levels of Lng108 in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in Lng108 levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of Lng108 is associated with a cancer which is regressing or in remission.

Further provided are methods of designing new therapeutic agents targeted to Lng108 for use in imaging and treating cancer. For example, in one embodiment, therapeutic agents such as antibodies targeted against Lng108 or fragments of such antibodies can be used to detect or image localization of Lng108 in a patient for the purpose of detecting or diagnosing a disease or condition. Such antibodies can be polyclonal, monoclonal, or omniconal or prepared by molecular biology techniques. The term "antibody", as used herein and throughout the instant specification is also meant to include aptamers and

-7-

single-stranded oligonucleotides such as those derived from an *in vitro* evolution protocol referred to as SELEX and well known to those skilled in the art. Antibodies can be labeled with a variety of detectable labels including, but not limited to, radioisotopes and paramagnetic metals.

Therapeutics agents such as small molecules and antibodies which decrease the concentration and/or activity of Lng108 can also be used in the treatment of diseases characterized by expression of Lng108. Such agents can be readily

identified in accordance with the teachings herein.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The present invention relates to diagnostic assays and methods, both quantitative and qualitative for detecting, diagnosing, monitoring, staging, prognosticating, *in vivo* imaging and treating cancers by comparing levels of Lng108 with those of Lng108 in a normal human control. Lng108 refers, among other things, to native proteins expressed by the gene comprising the polynucleotide sequence of SEQ ID

NO:1 or 2. The deduced amino acid sequence of a polypeptide encoded thereby is depicted in SEQ ID NO:3. By "Lng108" it is also meant herein polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1 or 2, but which still encode the same protein. In the alternative,

-8-

what is meant by Lng108 as used herein, means the native mRNA encoded by the gene comprising SEQ ID NO:1 or 2 or it can refer to the actual gene comprising SEQ ID NO:1 or 2, or levels of a polynucleotide which is capable of

5 hybridizing under stringent conditions to the antisense sequence of SEQ ID NO:1 or 2. Such levels are preferably measured in at least one of, cells, tissues and/or bodily fluids, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in

10 accordance with the invention for diagnosing over-expression of Lng108 protein compared to normal control bodily fluids, cells, or tissue samples may be used to diagnose the presence of cancers, including lung cancer. Lng108 may be measured alone in the methods of the

15 invention, or, more preferably, in combination with other diagnostic markers for cancer. Thus, it is preferred that the methods of the present invention be employed in combination with measurement of the levels of other cancer markers as well as Lng108. Other cancer markers, in

20 addition to Lng108, useful in the present invention will depend on the cancer being tested and are known to those of skill in the art.

Detection of Lng108 is particularly useful in lung cancer. However, this marker is also useful in the

25 diagnosis, prognosis, staging, imaging and treatment of other types of cancer.

Diagnostic Assays

The present invention provides methods for diagnosing the presence of cancer, including lung cancer, by analyzing

30 for changes in levels of Lng108 in cells, tissues or bodily fluids compared with levels of Lng108 in cells, tissues or bodily fluids of preferably the same type from a normal human control, wherein an increase in levels of Lng108 in the patient versus the normal human control is associated

35 with the presence of cancer.

-9-

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the patient being tested has cancer is one in which cells, tissues, or bodily fluid levels of a cancer marker, such as Lng108, are at least two times higher, and most preferable are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal human control.

The present invention also provides a method of diagnosing metastatic cancer, including metastatic lung cancer, in a patient having a cancer which has not yet metastasized. In the method of the present invention, a human cancer patient suspected of having cancer which may have metastasized (but which was not previously known to have metastasized) is identified. This is accomplished by a variety of means known to those of skill in the art.

In the present invention, determining the presence of Lng108 in cells, tissues, or bodily fluid, is particularly useful for discriminating between cancers which have not metastasized and cancers which have metastasized. Existing techniques have difficulty discriminating between a cancer which has metastasized and a cancer which has not metastasized and proper treatment selection is often dependent upon such knowledge.

In the present invention, one of the cancer marker levels measured in cells, tissues, or bodily fluid of a human patient is Lng108. Levels in the human patient are compared with levels of Lng108 in preferably the same cells, tissue, or bodily fluid type of a normal human control. That is, if the cancer marker being observed is Lng108 in serum, this level is preferably compared with the level of Lng108 in serum of a normal human control. An increase in Lng108 in the human patient versus the normal human control is associated with a cancer which has metastasized.

-10-

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the cancer in the patient being tested or monitored has metastasized is one in which cells, tissues, 5 or bodily fluid levels of a cancer marker, such as Lng108, are at least two times higher, and more preferably are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal human control.

Normal human control as used herein includes a human 10 patient without cancer and/or non cancerous samples from the patient; in the methods for diagnosing or monitoring for metastasis, normal human control may preferably also include samples from a human patient that is determined by reliable methods to have a cancer such as lung cancer which 15 has not metastasized.

Staging

The invention also provides a method of staging cancers in a human patient. The method comprises identifying a human patient having cancer and analyzing a 20 sample of cells, tissues, or bodily fluid from such patient for Lng108. The measured Lng108 levels are then compared to levels of Lng108 in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in Lng108 levels in the human patient versus the 25 normal human control is associated with a cancer which is progressing and a decrease in the levels of Lng108 is associated with a cancer which is regressing or in remission.

Monitoring

30 Further provided is a method of monitoring cancer in a human patient for the onset of metastasis. The method comprises identifying a human patient having cancer that is not known to have metastasized; periodically analyzing cells, tissues, or bodily fluid from such patient for 35 Lng108; and comparing the Lng108 levels in such cells,

-11-

tissue, or bodily fluid with levels of Lng108 in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in Lng108 levels in the patient versus the normal human control is associated with
5 a cancer which has metastasized.

Further provided by this invention is a method of monitoring the change in stage of a cancer. The method comprises identifying a human patient having cancer; periodically analyzing cells, tissues, or bodily fluid from
10 such patient for Lng108; and comparing the Lng108 levels in such cells, tissue, or bodily fluid with levels of Lng108 in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in Lng108 levels in the patient versus the normal human control is
15 associated with a cancer which is progressing in stage and a decrease in the levels of Lng108 is associated with a cancer which is regressing in stage or in remission.

Monitoring such patients for onset of metastasis is periodic and preferably done on a quarterly basis.
20 However, this may be performed more or less frequent depending on the cancer, the particular patient, and the stage of the cancer.

Prognostic Testing and Clinical Trial Monitoring

The methods described herein can further be utilized
25 as prognostic assays to identify subjects having or at risk of developing a disease or disorder associated with increased levels of Lng108. The present invention provides a method in which a test sample is obtained from a human patient and Lng108 is detected. The presence of higher
30 Lng108 levels as compared to normal human controls is diagnostic for the human patient being at risk for developing cancer, particularly lung cancer.

The effectiveness of therapeutic agents to decrease expression or activity of Lng108 can also be monitored by
35 analyzing levels of expression of Lng108 in a human patient

-12-

in clinical trials or in *in vitro* screening assays such as in human cells. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the human patient, or cells as the case may be, to the agent being tested.

Detection of genetic lesions or mutations

The methods of the present invention can also be used to detect genetic lesions or mutations in Lng108, thereby determining if a human with the genetic lesion is at risk for cancer or has cancer, particularly lung cancer. Genetic lesions can be detected, for example, by ascertaining the existence of a deletion and/or addition and/or substitution of one or more nucleotides from Lng108, a chromosomal rearrangement of Lng108, aberrant modification of Lng108 (such as of the methylation pattern of the genomic DNA), the presence of a non-wild type splicing pattern of a mRNA transcript of Lng108, allelic loss of Lng108, and/or inappropriate post-translational modification of Lng108 protein. Methods to detect such lesions in Lng108 are known to those of skill in the art.

Assay Techniques

Assay techniques that can be used to determine levels of gene expression, such as Lng108 of the present invention, in a sample derived from a human are well-known to those of skill in the art. Such assay methods include radioimmunoassays, reverse transcriptase PCR (RT-PCR) assays, immunohistochemistry assays, *in situ* hybridization assays, competitive-binding assays, Western Blot analyses, ELISA assays and proteomic approaches. Among these, ELISAs are frequently preferred to diagnose a gene's expressed protein in biological fluids.

An ELISA assay initially comprises preparing an antibody, if not readily available from a commercial source, specific to Lng108, preferably a monoclonal antibody. In addition a reporter antibody generally is

-13-

prepared which binds specifically to Lng108. The reporter antibody is attached to a detectable reagent such as a radioactive, fluorescent or enzymatic reagent, for example horseradish peroxidase enzyme or alkaline phosphatase.

5 To carry out the ELISA, antibody specific to Lng108 is incubated on a solid support, e.g., a polystyrene dish, that binds the antibody. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein such as bovine serum albumin. Next, the sample to
10 be analyzed is incubated in the dish, during which time Lng108 binds to the specific antibody attached to the polystyrene dish. Unbound sample is washed out with buffer. A reporter antibody specifically directed to Lng108 and linked to a detectable reagent such as horseradish
15 peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to Lng108. Unattached reporter antibody is then washed out. Reagents for peroxidase activity, including a colorimetric substrate are then added to the dish. Immobilized
20 peroxidase, linked to Lng108 antibodies, produces a colored reaction product. The amount of color developed in a given time period is proportional to the amount of Lng108 protein present in the sample. Quantitative results typically are obtained by reference to a standard curve.

25 A competition assay can also be employed wherein antibodies specific to Lng108 are attached to a solid support and labeled Lng108 and a sample derived from the patient or human control are passed over the solid support. The amount of label detected which is attached to the solid
30 support can be correlated to a quantity of Lng108 in the sample.

Using all or a portion of the nucleic acid sequence for Lng108 as a hybridization probe, nucleic acid methods can also be used to detect Lng108 mRNA as a marker for
35 cancer, including lung cancer. Polymerase chain reaction

-14-

(PCR) and other nucleic acid methods, such as ligase chain reaction (LCR) and nucleic acid sequence based amplification (NASABA), can be used to detect malignant cells for diagnosis and monitoring of various malignancies.

5 For example, reverse-transcriptase PCR (RT-PCR) is a powerful technique which can be used to detect the presence of a specific mRNA population in a complex mixture of thousands of other mRNA species. In RT-PCR, an mRNA species is first reverse transcribed to complementary DNA
10 (cDNA) with use of the enzyme reverse transcriptase; the cDNA is then amplified as in a standard PCR reaction. RT-PCR can thus reveal by amplification the presence of a single species of mRNA. Accordingly, if the mRNA is highly specific for the cell that produces it, RT-PCR can be used
15 to identify the presence of a specific type of cell.

Hybridization to clones or oligonucleotides arrayed on a solid support (i.e., gridding) can be used to both detect the expression of and quantitate the level of expression of that gene. In this approach, a cDNA encoding the Lng108
20 gene is fixed to a substrate. The substrate may be of any suitable type including but not limited to glass, nitrocellulose, nylon or plastic. At least a portion of the DNA encoding the Lng108 gene is attached to the substrate and then incubated with the analyte, which may be
25 RNA or a complementary DNA (cDNA) copy of the RNA, isolated from the tissue of interest. Hybridization between the substrate bound DNA and the analyte can be detected and quantitated by several means including but not limited to radioactive labeling or fluorescence labeling of the
30 analyte or a secondary molecule designed to detect the hybrid. Quantitation of the level of gene expression can be done by comparison of the intensity of the signal from the analyte compared with that determined from known standards. The standards can be obtained by *in vitro*
35 transcription of the target gene, quantitating the yield,

-15-

and then using that material to generate a standard curve.

Of the proteomic approaches, 2D electrophoresis is a technique well known to those in the art. Isolation of individual proteins from a sample such as serum is
5 accomplished using sequential separation of proteins by different characteristics usually on polyacrylamide gels. First, proteins are separated by size using an electric current. The current acts uniformly on all proteins, so smaller proteins move farther on the gel than larger
10 proteins. The second dimension applies a current perpendicular to the first and separates proteins not on the basis of size but on the specific electric charge carried by each protein. Since no two proteins with different sequences are identical on the basis of both size
15 and charge, the result of a 2D separation is a square gel in which each protein occupies a unique spot. Analysis of the spots with chemical or antibody probes, or subsequent protein microsequencing can reveal the relative abundance of a given protein and the identity of the proteins in the
20 sample.

The above tests can be carried out on samples derived from a variety cells, bodily fluids and/or tissue extracts (homogenates or solubilized tissue) obtained from the patient including tissue biopsy and autopsy material.
25 Bodily fluids useful in the present invention include blood, urine, saliva, or any other bodily secretion or derivative thereof. Blood can include whole blood, plasma, serum, or any derivative of blood.

In Vivo Targeting of Lng108/Cancer Therapy

30 Identification of Lng108 is also useful in the rational design of new therapeutics for imaging and treating cancers, and in particular lung cancer. For example, in one embodiment, antibodies which specifically bind to Lng108 can be raised and used *in vivo* in patients
35 suspected of suffering from cancer. Antibodies which

-16-

specifically bind a Lngl08 can be injected into a patient suspected of having cancer for diagnostic and/or therapeutic purposes. Thus, another aspect of the present invention provides for a method for preventing the onset and treatment of lung cancer in a human patient in need of such treatment by administering to the patient an effective amount of an antibody. By "effective amount" it is meant the amount or concentration of antibody needed to bind to the target antigens expressed on the tumor to cause tumor shrinkage for surgical removal, or disappearance of the tumor. The binding of the antibody to Lngl08 is believed to cause the death of the cancer cell expressing such Lngl08. The preparation and use of antibodies for *in vivo* diagnosis is well known in the art. For example, antibody-chelators labeled with Indium-111 have been described for use in the radioimmunoscentographic imaging of carcinoembryonic antigen expressing tumors (Sumerdon et al. Nucl. Med. Biol. 1990 17:247-254). In particular, these antibody-chelators have been used in detecting tumors in patients suspected of having recurrent colorectal cancer (Griffin et al. J. Clin. Onc. 1991 9:631-640). Antibodies with paramagnetic ions as labels for use in magnetic resonance imaging have also been described (Lauffer, R.B. Magnetic Resonance in Medicine 1991 22:339-342).

Antibodies directed against Lngl08 can be used in a similar manner. Labeled antibodies which specifically bind Lngl08 can be injected into patients suspected of having cancer for the purpose of diagnosing or staging of the disease status of the patient. The label used will be selected in accordance with the imaging modality to be used. For example, radioactive labels such as Indium-111, Technetium-99m or Iodine-131 can be used for planar scans or single photon emission computed tomography (SPECT). Positron emitting labels such as Fluorine-19 can be used in positron emission tomography. Paramagnetic ions such as Gadlinium

-17-

(III) or Manganese (II) can be used in magnetic resonance imaging (MRI). Localization of the label permits determination of the spread of the cancer. The amount of label within an organ or tissue also allows determination of the presence or absence of cancer in that organ or tissue.

Antibodies which can be used in *in vivo* methods include polyclonal, monoclonal and omniclonal antibodies and antibodies prepared via molecular biology techniques. Antibody fragments and aptamers and single-stranded oligonucleotides such as those derived from an *in vitro* evolution protocol referred to as SELEX and well known to those skilled in the art can also be used.

Screening Assays

The present invention also provides methods for identifying modulators which bind to Lng108 protein or have a modulatory effect on the expression or activity of Lng108 protein. Modulators which decrease the expression or activity of Lng108 protein are believed to be useful in treating cancer, particularly lung cancer. Such screening assays are known to those of skill in the art and include, without limitation, cell-based assays and cell free assays.

Small molecules predicted via computer imaging to specifically bind to regions of Lng108 can also be designed, synthesized and tested for use in the imaging and treatment of cancer. Further, libraries of molecules can be screened for potential anticancer agents by assessing the ability of the molecule to bind to Lng108. Molecules identified in the library as being capable of binding to Lng108 are key candidates for further evaluation for use in the treatment of cancer, particularly lung cancer. In a preferred embodiment, these molecules will downregulate expression and/or activity of Lng108 in cells.

Adoptive Immunotherapy and Vaccines

Adoptive immunotherapy of cancer refers to a

-18-

therapeutic approach in which immune cells with an antitumor reactivity are administered to a tumor-bearing host, with the aim that the cells mediate either directly or indirectly, the regression of an established tumor.

- 5 Transfusion of lymphocytes, particularly T lymphocytes, falls into this category and investigators at the National Cancer Institute (NCI) have used autologous reinfusion of peripheral blood lymphocytes or tumor-infiltrating lymphocytes (TIL), T cell cultures from biopsies of
10 subcutaneous lymph nodes, to treat several human cancers (Rosenberg, S. A., U.S. Patent No. 4,690,914, issued Sep. 1, 1987; Rosenberg, S. A., et al., 1988, N. England J. Med. 319:1676-1680).

The present invention relates to compositions and
15 methods of adoptive immunotherapy for the prevention and/or treatment of primary and metastatic cancer in humans using macrophages sensitized to the antigenic Lng108 molecules, with or without non-covalent complexes of heat shock protein (hsp). Antigenicity or immunogenicity of Lng108 is
20 readily confirmed by the ability of the Lng108 protein or a fragment thereof to raise antibodies or educate naive effector cells, which in turn lyse target cells expressing the antigen (or epitope).

Cancer cells are, by definition, abnormal and contain
25 proteins which should be recognized by the immune system as foreign since they are not present in normal tissues. However, the immune system often seems to ignore this abnormality and fails to attack tumors. The foreign Lng108 proteins that are produced by the cancer cells can be used
30 to reveal their presence. The Lng108 is broken into short fragments, called tumor antigens, which are displayed on the surface of the cell. These tumor antigens are held or presented on the cell surface by molecules called MHC, of which there are two types: class I and II. Tumor antigens
35 in association with MHC class I molecules are recognized by

-19-

cytotoxic T cells while antigen-MHC class II complexes are recognized by a second subset of T cells called helper cells. These cells secrete cytokines which slow or stop tumor growth and help another type of white blood cell, B cells, to make antibodies against the tumor cells.

In adoptive immunotherapy, T cells or other antigen presenting cells (APCs) are stimulated outside the body (*ex vivo*), using the tumor specific Lng108 antigen. The stimulated cells are then reinfused into the patient where they attack the cancerous cells. Research has shown that using both cytotoxic and helper T cells is far more effective than using either subset alone. Additionally, the Lng108 antigen may be complexed with heat shock proteins to stimulate the APCs as described in U.S. Patent No. 5,985,270.

The APCs can be selected from among those antigen presenting cells known in the art including, but not limited to, macrophages, dendritic cells, B lymphocytes, and a combination thereof, and are preferably macrophages. In a preferred use, wherein cells are autologous to the individual, autologous immune cells such as lymphocytes, macrophages or other APCs are used to circumvent the issue of whom to select as the donor of the immune cells for adoptive transfer. Another problem circumvented by use of autologous immune cells is graft versus host disease which can be fatal if unsuccessfully treated.

In adoptive immunotherapy with gene therapy, DNA of the Lng108 can be introduced into effector cells similarly as in conventional gene therapy. This can enhance the cytotoxicity of the effector cells to tumor cells as they have been manipulated to produce the antigenic protein resulting in improvement of the adoptive immunotherapy.

Lng108 antigens of this invention are also useful as components of cancer vaccines. The vaccine comprises an immunogenically stimulatory amount of a Lng108 antigen.

-20-

Immunogenically stimulatory amount refers to that amount of antigen that is able to invoke the desired immune response in the recipient for the amelioration, or treatment of cancer, particularly lung cancer. Effective amounts may be
5 determined empirically by standard procedures well known to those skilled in the art.

The Lng108 antigen may be provided in any one of a number of vaccine formulations which are designed to induce the desired type of immune response, e.g., antibody and/or
10 cell mediated. Such formulations are known in the art and include, but are not limited to, formulations such as those described in U.S. Patent 5,585,103. Vaccine formulations of the present invention used to stimulate immune responses can also include pharmaceutically acceptable adjuvants.

15 **EXAMPLE**

The present invention is further described by the following example. The example is provided solely to illustrate the invention by reference to specific
embodiments. This exemplification, while illustrating
20 certain specific aspects of the invention, does not portray the limitations or circumscribe the scope of the disclosed invention.

Experiments described herein were carried out using standard techniques, which are well known and routine to
25 those of skill in the art, except where otherwise described in detail. Routine molecular biology techniques were carried out as described in standard laboratory manuals, such as Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold
30 Spring Harbor, N.Y. (1989).

Relative Quantitation of Gene Expression

Real-Time quantitative PCR with fluorescent Taqman probes is a quantitation detection system utilizing the 5'-
3' nuclease activity of Taq DNA polymerase. The method
35 uses an internal fluorescent oligonucleotide probe (Taqman)

-21-

labeled with a 5' reporter dye and a downstream, 3' quencher dye. During PCR, the 5'-3' nuclease activity of Taq DNA polymerase releases the reporter, whose fluorescence can then be detected by the laser detector of the Model 7700 Sequence Detection System (PE Applied Biosystems, Foster City, CA, USA).

Amplification of an endogenous control was used to standardize the amount of sample RNA added to the reaction and normalize for Reverse Transcriptase (RT) efficiency. Either cyclophilin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) or 18S ribosomal RNA (rRNA) was used as this endogenous control. To calculate relative quantitation between all the samples studied, the target RNA levels for one sample were used as the basis for comparative results (calibrator). Quantitation relative to the "calibrator" can be obtained using the standard curve method or the comparative method (User Bulletin #2: ABI PRISM 7700 Sequence Detection System).

The tissue distribution, and the level of the target gene for every example in normal and cancer tissue were determined. Total RNA was extracted from normal tissues, cancer tissues, and from cancers and the corresponding matched adjacent tissues. Subsequently, first strand cDNA was prepared with reverse transcriptase and the polymerase chain reaction was done using primers and Taqman probe specific to each target gene. The results are analyzed using the ABI PRISM 7700 Sequence Detector. The absolute numbers are relative levels of expression of the target gene in a particular tissue compared to the calibrator tissue.

Primers used for expression analysis include:

5' TCTAGGTCAGCCCCGAATC 3' (SEQ ID NO:4); and

5' CCTCCAATTCCCCCTTAAACTT 3' (SEQ ID NO:5).

The absolute numbers depicted in Table 1 are relative levels of expression of Lng108 (also referred to as Clone

-22-

ID 954287; Gene ID 21300) in 12 normal different tissues. All the values are compared to normal muscle (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from
 5 different individuals.

Table 1: Relative Levels of Lng108 Expression in Pooled Samples

	TISSUE	NORMAL
10	Brain	0.57
	Heart	1.63
	Kidney	9.55
	Liver	0.38
	Lung	53.46
15	Mammary Gland	13.00
	Muscle	1.00
	Prostate	1.69
	Small Intestine	0.80
	Testis	0.56
20	Thymus	1.06
	Uterus	4.88

The relative levels of expression in Table 1 show that Lng108 mRNA is expressed in all 12 tissue types analyzed. The expression level of Lng108 is relatively higher in lung and is lower in brain, liver, small intestine and testis
 25 and is medium in kidney, mammary gland and uterus. These results demonstrate that Lng108 mRNA expression is not restricted to lung tissue but is expressed broadly in all tissue types analyzed.

The absolute numbers in Table 1 were obtained
 30 analyzing pools of samples of a particular tissue from different individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 2.

The absolute numbers depicted in Table 2 are relative
 35 levels of expression of Lng108 in 48 pairs of matching samples and 2 cancer and 2 normal/normal adjacent tissues of ovary. All the values are compared to normal muscle (calibrator). A matching pair is formed by mRNA from the

-23-

cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 2: Relative Levels of Lngl08 Expression in Individual Samples

5

10

15

20

Sample ID	Cancer Type	Tissue	Cancer	Matching Normal Adjacent
LngAC82	Adenocarcinoma	Lung 1	29.75	28.15
Lng60XL	Adenocarcinoma	Lung 2	28.9	5.3
LngAC66	Adenocarcinoma	Lung 3	5.01	5.50
LngAC69	Adenocarcinoma	Lung 4	58.28	15.19
LngAC88	Adenocarcinoma	Lung 5	90.20	111.0
LngAC13	Adenocarcinoma	Lung 6	18.32	0.00
LngSQ9X	Squamous cell carcinoma	Lung 7	57.88	9.09
LngQ45	Squamous cell carcinoma	Lung 8	31	76
LngSQ56	Squamous cell carcinoma	Lung 9	56	65
LngSQ32	Squamous cell carcinoma	Lung 10	3821.7	218.3
LngSQ79	Squamous cell carcinoma	Lung 11	574.04	467.88
LngC20X	Squamous cell carcinoma	Lung 12	0.7	0.4
Lng47XQ	Squamous cell carcinoma	Lung 13	204	0.5
LngSQ44	Squamous cell carcinoma	Lung 14	8.70	61.20
LngBR94	Squamous cell carcinoma	Lung 15	85.0	0.0
Lng90X	Squamous cell carcinoma	Lung 16	12.7	7.3
LngLC71	Large cell carcinoma	Lung 17	82.14	71.26

-24-

	Lng LC109	Large cell carcinoma	Lung 18	94.35	348.50
	Lng75XC	Metastatic from bone cancer	Lung 19	1	3
	LngMT67	Metastatic from renal cell cancer	Lung 20	590.18	20.04
5	LngMT71	Metastatic from melanoma	Lung 21	32.90	18.06
	Bld32XK		Bladder 1	17.5	4.6
	Bld46XK		Bladder 2	3.4	5.8
	ClnAS67		Colon 1	28.4	0.1
	ClnC9XR		Colon 2	29	10
10	ClnTX67		Colon 3	78	2
	End28XA		Endometrium 1	49.2	35.4
	End12XA		Endometrium 2	13	13
	Kid 106XD		Kidney 1	16.6	52.2
15	Kid 107XD		Kidney 2	1992.0	61.0
	Kid 109XD		Kidney 3	641	53
	Liv94XA		Liver 1	30.0	1.6
20	Liv15XA		Liver 2	5	3
	MamA06X		Liver 3	20.9	1.1
	Mam B011X		Mammary gland 1	46.7	0.2
	Mam12X		Mammary gland 2	80	97
25	Ovr103X		Ovary 1	44.2	0.9
	Ovr 10050		Ovary 2	40	
	Ovr1028		Ovary 3	136	

-25-

	Ovr18GA		Ovary 4		116
	Ovr206I		Ovary 5		5
	Pan71XL		Pancreas 1	0.5	0.4
	Pan77X		Pancreas 2	21	7
5	Pro20XB		Prostate 1	2.9	15.9
	Pro12B		Prostate 2	11	2
	Pro13XB		Prostate 3	0.6	10
	SmIH89		Small Intestine 1	28	6
	StoAC44		Stomach 1	8	24
10	Tst39X		Testis 1	184.8	1.8
	Utr 135XO		Uterus 1	88.0	138.6
	Utr 141XO		Uterus 2	110	65
15	Utr23XU		Uterus 3	58	41

0= Negative

In the analysis of matching samples, the higher levels of expression were in lung tissue. In addition to the expression in lung, Lngl08 was also expressed in all other 14 tissue types tested. These results confirmed that Lngl08 is expressed higher in lung but also is expressed in other tissue types analyzed and is consistent with the results obtained with the panel of normal pooled samples (Table 1).

Furthermore, the level of mRNA expression was compared in cancer samples and the isogenic normal adjacent tissue from the same individual. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 2 shows overexpression of Lngl08 in 14 out of 21 (67%) lung cancer tissues compared with their respective normal adjacent. Overexpression of Lngl08 was also found in other cancer samples compared to

-26-

the normal adjacent tissues (bladder, colon, endometrium, kidney, liver, mammary, ovary, prostate, small intestine, testis and uterus). Overall, these results show overexpression of Lng108 in 36 out of 48 (75%) cancer

5 tissues tested compared to the normal adjacent.

Thus, the mRNA expression in many different tissue types, plus the observed overexpression in 75% of all the cancer matching samples tested is indicative of Lng108 being a lung cancer diagnostic marker and a general cancer

10 diagnostic marker.

-27-

What is claimed is:

1. A method for diagnosing the presence of cancer in a patient comprising:

(a) determining levels of Lng108 in cells, tissues or
5 bodily fluids in a patient; and

(b) comparing the determined levels of Lng108 with levels of Lng108 in cells, tissues or bodily fluids from a normal human control, wherein a change in determined levels of Lng108 in said patient versus normal human control is
10 associated with the presence of cancer.

2. A method of diagnosing metastases of cancer in a patient comprising:

(a) identifying a patient having cancer that is not known to have metastasized;

15 (b) determining Lng108 levels in a sample of cells, tissues, or bodily fluid from said patient; and

(c) comparing the determined Lng108 levels with levels of Lng108 in cells, tissue, or bodily fluid of a normal human control, wherein an increase in determined Lng108
20 levels in the patient versus the normal human control is associated with a cancer which has metastasized.

3. A method of staging cancer in a patient having cancer comprising:

(a) identifying a patient having cancer;

25 (b) determining Lng108 levels in a sample of cells, tissue, or bodily fluid from said patient; and

(c) comparing determined Lng108 levels with levels of Lng108 in cells, tissues, or bodily fluid of a normal human control, wherein an increase in determined Lng108 levels in
30 said patient versus the normal human control is associated with a cancer which is progressing and a decrease in the determined Lng108 levels is associated with a cancer which is regressing or in remission.

4. A method of monitoring cancer in a patient for
35 the onset of metastasis comprising:

-28-

(a) identifying a patient having cancer that is not known to have metastasized;

(b) periodically determining levels of Lngl08 in samples of cells, tissues, or bodily fluid from said patient; and

(c) comparing the periodically determined Lngl08 levels with levels of Lngl08 in cells, tissues, or bodily fluid of a normal human control, wherein an increase in any one of the periodically determined Lngl08 levels in the patient versus the normal human control is associated with a cancer which has metastasized.

5. A method of monitoring a change in stage of cancer in a patient comprising:

(a) identifying a patient having cancer;

(b) periodically determining levels of Lngl08 in cells, tissues, or bodily fluid from said patient; and

(c) comparing the periodically determined Lngl08 levels with levels of Lngl08 in cells, tissues, or bodily fluid of a normal human control, wherein an increase in any one of the periodically determined Lngl08 levels in the patient versus the normal human control is associated with a cancer which is progressing in stage and a decrease is associated with a cancer which is regressing in stage or in remission.

6. A method of identifying potential therapeutic agents for use in imaging and treating cancer comprising screening molecules for an ability to bind to or decrease expression of Lngl08 wherein the ability of a molecule to bind to Lngl08 or decrease expression of Lngl08 is indicative of the molecule being useful in imaging and treating cancer.

7. A method of imaging cancer in a patient comprising administering to the patient an antibody which specifically binds to Lngl08.

8. The method of claim 7 wherein said antibody is

-29-

labeled with paramagnetic ions or a radioisotope.

9. A method of treating cancer in a patient comprising administering to the patient an antibody which specifically binds to Lng108.

5 10. The method of claim 9 wherein the antibody is conjugated to a cytotoxic agent.

11. A method of treating cancer in a patient comprising administering to the patient a molecule which downregulates expression or activity of Lng108.

10 12. The method of claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11 wherein the cancer is lung cancer.

13. A method of inducing an immune response against a target cell expressing Lng108 comprising delivering to a human patient an immunogenically stimulatory amount of a
15 Lng108 protein so that an immune response is mounted against the target cell.

14. A vaccine for treating cancer comprising an immunogenically stimulating amount of Lng108.

SEQUENCE LISTING

<110> Recipon, Herve

Macina, Roberto A.

Chen, Sei-Yu

Sun, Yongming

DIADEXUS, INC.

<120> A Novel Method of Diagnosing, Monitoring, Staging,
Imaging and Treating Cancer

<130> DEX-0088

<140>

<141>

<150> 60/163,444

<151> 1999-11-04

<160> 5

<170> PatentIn Ver. 2.1

<210> 1

<211> 503

<212> DNA

<213> Homo sapiens

<400> 1

```

ctatatatgt atctacaata catatatcta cacatacaga aagaagcagt tctcacaatg 60
ttgctagttt tttgcttctc tttcccccac cctactccct ccaattcccc cttaaacttc 120
caaagcttcg tcttggtgtt gctgcagagt gattcggggg ctgacctaga ccagtttgca 180
tgattcttct cttgtgattt ggttgcaatt tagacatttt tgtgccatta tatttgcat 240
atgtatttat aatttaaatg atatttaggt ttttggtga gtactggaat aaacagtga 300
catatctggt atatgtcatt atttattggt aaattacatt ttttaagctcc atgtgcatat 360
aaaggttatg aaacatatca tggtaatgac agatgcaagt tattttattt gcttattttt 420
ataattaaag atgccatagc ataatatgaa gcctttggtg aattccttct aagataaaaa 480
taataataaa gtgttacgtt tta                                     503

```

<210> 2

<211> 3762

<212> DNA

<213> Homo sapiens

<400> 2

```

ggtggcagca gcagcatcac acgtaacaac aacaaaaaaaa aatcctcatc aaatcctcac 60
ctaagctttc agtgtatcca gatccacatc ttcactcaag ccaggagagg gaaagaggaa 120

```

```

aggggggcag gaaaaaaaaa aaacccaaca acttagcgga aacttctcag agaattgctcc 180
aaaactcagc agtgcttctg gtgtggtgga tcagtgtctt tgcaacccat gaggcggagc 240
agaatgactc tgtgagcccc aggaaatccc gaggcgggc tcaaaactca gctgaagtgg 300
ttcgttgcc tcaacagtgt ctacaggtcg gctgcggggc ttttgcatgc ctggaaaact 360
ccacctgtga cacagatggg atgtatgaca tctgtaaatc cttcttgtag agcgtgctta 420
aatgtgacac tcagggaaaa gcattcgtca aagagagctt aaaatgcac gccaacgggg 480
tcacctccaa ggtcttccct gccattcgga ggtgctccac tttccaaagg atgattgctg 540
aggtgcagga agagtgtctac agcaagctga atgtgtgcag catcgccaag cggaaccctg 600
aagccatcac tgaggtcgtc cagctgcccc atcacttctc caacagatac tataacagac 660
ttgtccgaag cctgctggaa tgtgatgaag acacagtcag cacaatcaga gacagcctga 720
tggagaaaat tgggcctaac atggccagcc tctccacat cctgcagaca gaccactgtg 780
cccaaacaca cccacgagct gacttcaaca ggagacgcac caatgagccg cagaagctga 840
aagtctcct caggaaacct cgaggtgagg aggactctcc ctccacatc aaacgcacat 900
cccattgagag tgcataacca gggagaggtt attcacaacc tcacaaact agtatcattt 960
taggggtgtt gacacaccag ttttgagtgt actgtgcctg gtttgatttt tttaaagtag 1020
ttcctatttt ctatccccct taaagaaaat tgcataaac taggcttctg taatcaatat 1080
cccaacattc tgcaatggca gcattcccac caacaaaatc catgtgacca ttctgcctct 1140
cctcaggaga aagtacctc ttttaccac ttcctctgcc atgttttcc cctgctcccc 1200
tgagaccacc cccaaacaca aaacattcat gtaactctcc agccattgta atttgaagat 1260
gtggatccct ttagaacggg tgccccagta gagttagctg ataaggaaac tttatttaa 1320
tgcattgtct aaatgtctat aaagatgtta aatggaattc gtgttatgaa tctgtgctgg 1380
ccatggacga atatgaatgt cacatttgaa ttcttgatct ctaatgagct agtgtcttat 1440
ggtcttgatc ctccaatgtc taattttctt tccgacacat ttaccaaatt gcttgagcct 1500
ggctgtccaa ccagactttg agcctgcatc ttcttgcatc taatgaaaaa caaaaagcta 1560
acattcttac gtactgtaac tgctcagagc tttaaaagta tctttaacaa ttgtcttaa 1620
accagagaat cttaaggtct aactgtggaa tataaatagc tgaaaactaa tgtactgtac 1680
ataaattcca gaggactctg cttaaacaaa gcagtatata ataactttat tgcataataga 1740
tttagttttg taacttagct ttattttctt tttcctggga atggaataac tatctcactt 1800
ccagatatcc acataaatgc tcttgtggc cttttttata actaaggggg tagaagtagt 1860
tttaattcaa catcaaaact taagatgggc ctgtatgaga caggaaaaac caacaggttt 1920
atctgaagga cccaggttaa gatgttaatc tcccagccca cctcaacca gaggtactc 1980
ttgacttaga cctatactga aagatctctg tcacatccaa ctggaaattc caggaaacca 2040
aaagagcacc ctatgggctt ggaccactta cagtgtgata aggcctacta tacattagga 2100
agtggcagtt ctttactcgt cccctttcat cgggtgcctg tactctggca aatgatgatg 2160
gggtgggaga ctttccatta aatcaatcag gaatgagtca atcagccttt aggtcttttag 2220
tccgggggac ttggggctga gagagtataa ataaccctgg gctgtccagc cttaatagac 2280
ttctcttaca ttttctcct gtagcacgct gcctgccaaa gtagtcctgg cagctggacc 2340
atctctgtag gatcgtaaaa aaatagaaan nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn 2400
nnnnnctggt ggttgatcat ttctgccatg atgtttacaa gatggcgacc accaangtca 2460
aacgactaac ctatctatga acaacagtag tttctcaggg tcaactgcct tgaacccaac 2520
agtcctttat gagcgtcact gccaccaaa ggtcaatgtc aagagaggaa gagaggagg 2580
aggggtagga ctgcaggggc cactccaaac tcgcttaggt agaaactatt ggtgcttgac 2640
tctcactagg ctaaactcaa gatttgacca aatcgagtga tagggatcct ggtgggagga 2700
gagagggcac atctccagaa aaatgaaaag caatacaact ttaccataaa gcctttaaa 2760
ccagtaacgt gctgctcaag gaccaagagc aattnnnnnn nnnnnnnnnn nnnnnnnnnn 2820
nnnncaaaca ttgctgctt tgtccccaca cagcctctaa gcgtgctgac atcagattgt 2880
taagggcatt ttataactca gaactgtccc atccccaggt ccccaaactt atggacactg 2940
ccttagcctc ttggaaatca ggtagaccat attctaagtt agactcttcc cctccctccc 3000

```



```

acacttccca cccccaggca aggetgactt ctctgaatca gaaaagctat taaagtttgt 3060
gtgttggtgc ctttttgcaa acccaactaa gccaggaccc caatgcgaca agtagttcat 3120
gagtattcct agcaaatttc tctctttctt cagttcagta gatttccttt tttcttntct 3180
ntnttttttt ntttttttgg ctgtgacctc ttcaaaccgt ggtaccccc cttttctccc 3240
cacgatgata tctatatatg tatctacaat acatatatct acacatacag aaagaagcag 3300
ttctcacaat gttgctagtt ttttgcttct ctttccccca ccctactccc tccaattccc 3360
ccttaaactt ccaaagcttc gtcttggtgt tgctgcagag tgattcgggg gctgacctag 3420
accagtttgc atgattcttc tcttggtgatt tggttgcact ttagacattt ttgtgccatt 3480
atatttgcac tatgtattta taatttaaag gatatttagg tttttggctg agtactggaa 3540
taaacagtga gcatatctgg tatatgtcat tatttattgt taaattacat ttttaagctc 3600
catgtgcata taaaggttat gaaacatata atggtaatac cagatgcaag ttattttatt 3660
tgcttatttt tataattaaa gatgccatag cataatatga agcctttggt gaattccttc 3720
taagataaaa ataataataa agtgttacgt tttattggtt tc 3762

```

<210> 3

<211> 247

<212> PRT

<213> Homo sapiens

<400> 3

```

Met Leu Gln Asn Ser Ala Val Leu Leu Val Leu Val Ile Ser Ala Ser
  1              5              10              15

```

```

Ala Thr His Glu Ala Glu Gln Asn Asp Ser Val Ser Pro Arg Lys Ser
      20              25              30

```

```

Arg Val Ala Ala Gln Asn Ser Ala Glu Val Val Arg Cys Leu Asn Ser
      35              40              45

```

```

Ala Leu Gln Val Gly Cys Gly Ala Phe Ala Cys Leu Glu Asn Ser Thr
      50              55              60

```

```

Cys Asp Thr Asp Gly Met Tyr Asp Ile Cys Lys Ser Phe Leu Tyr Ser
      65              70              75              80

```

```

Ala Ala Lys Phe Asp Thr Gln Gly Lys Ala Phe Val Lys Glu Ser Leu
      85              90              95

```

```

Lys Cys Ile Ala Asn Gly Val Thr Ser Lys Val Phe Leu Ala Ile Arg
      100             105             110

```

```

Arg Cys Ser Thr Phe Gln Arg Met Ile Ala Glu Val Gln Glu Glu Cys
      115             120             125

```

```

Tyr Ser Lys Leu Asn Val Cys Ser Ile Ala Lys Arg Asn Pro Glu Ala
      130             135             140

```

Ile Thr Glu Val Val Gln Leu Pro Asn His Phe Ser Asn Arg Tyr Tyr
145 150 155 160

Asn Arg Leu Val Arg Ser Leu Leu Glu Cys Asp Glu Asp Thr Val Ser
165 170 175

Thr Ile Arg Asp Ser Leu Met Glu Lys Ile Gly Pro Asn Met Ala Ser
180 185 190

Leu Phe His Ile Leu Gln Thr Asp His Cys Ala Gln Thr His Pro Arg
195 200 205

Ala Asp Phe Asn Arg Arg Arg Thr Asn Glu Pro Gln Lys Leu Lys Val
210 215 220

Leu Leu Arg Asn Leu Arg Gly Glu Glu Asp Ser Pro Ser His Ile Lys
225 230 235 240

Arg Thr Ser His Glu Ser Ala
245

<210> 4

<211> 20

<212> DNA

<213> Homo sapiens

<400> 4

tctaggtcag cccccgaatc

20

<210> 5

<211> 22

<212> DNA

<213> Homo sapiens

<400> 5

cctccaattc ccccttaaac tt

22

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/30482

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 39/395, 39/40, 39/42, 39/44, 39/00, 39/38, 38/00; C12N 5/06, 5/16
US CL : 424/138.1, 178.1, 184.1, 277.1; 435/330; 514/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/138.1, 178.1, 184.1, 277.1; 435/330; 514/12

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Continuation Sheet

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X/P	FUJIWARA, Y. et al. Assessment of stanniocalcin-1 mRNA as a molecular marker for micrometases of various human cancers. Int. J. Oncol. April 2000, Vol. 16, pages 799-804, especially page 801.	1-2
A,P		3-14

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

23 January 2001

Date of mailing of the international search report

20 MAR 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Bruce William Tedeschi

Telephone No. 703-306-4823

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/30482

Continuation of B. FIELDS SEARCHED Item3: BIOSIS, CAPLUS, WPIDS, USPATFULL, MEDLINE, CANCERLIT, WEST

search terms: stanniocalcin, Lng108, cancer, lung cancer, metastasis, antibody, vaccine